

The Constraint on FCNC Coupling of the Top Quark with a Gluon from ep - Collisions

A.A. Ashimova ¹

Moscow State University, 119992 Moscow, Russia

S.R. Slabospitsky ²

State Research Center

Institute for High Energy Physics,

Protvino, Moscow Region 142281, Russia

Abstract

Using the constraint on the single top production cross-section obtained at the HERA collider, $\sigma(ep \rightarrow e t X)$, we evaluate an upper limit on coupling constant of the anomalous top quark interaction with a gluon via flavor-changing neutral current: $|\kappa_{tgq}/\Lambda| \lesssim 0.4 \text{ TeV}^{-1}$, $\text{BR}(t \rightarrow gq) < 13\%$

¹E-mail: ashimova_aa@mail.ru

²E-mail: Sergey.Slabospitsky@ihep.ru

In different extensions of the Standard Model (SM) the anomalous interactions can lead to significant modification of the top quark production mechanisms and appearance of the top rare decays [1]. Thereby, the precision measurements of the top quark properties and its production mechanisms provide a possibility to obtain an information on new physics beyond the SM. Of special interest it is the study of the top quark anomalous interactions via Flavour-Changing Neutral Currents (FCNC).

In the SM there are no direct FCNC transitions $t \rightarrow u(c)$. Only "loop" contributions can make them possible. As a result such processes are strongly suppressed within the SM: $\text{BR}(t \rightarrow (\gamma, g, Z) + u(c)) < \mathcal{O}(10^{-10})$ [2]. However, various extensions of the SM could lead to a huge enhancement of FCNC processes [3, 4, 5].

Search for the anomalous FCNC interactions of the top quark were performed at the Tevatron [6], HERA [7, 8, 9], and LEP-2 [10] colliders. The present constraints on the top quark anomalous couplings in terms of branching ratio (BR) are presented in the Table below.

Table 1: *The present constraints on top quark anomalous couplings*

decay	BR	collider
$t \rightarrow q\gamma$	$< 0.7\%$	HERA [8, 9]
$t \rightarrow qZ$	$< 7\%$	LEP 2 [10]
$t \rightarrow qg$	$< 8\%$	Tevatron + th. [11]

Search for the top production at HERA collider in ep -collisions,

$$ep \rightarrow etX, \quad (1)$$

was done by the H1 [8] and ZEUS [9] collaborations at the center-mass energy of $\sqrt{s} = 318$ GeV. Obviously that only single top production is possible at this energy. In this case the SM dominant process is $ep \rightarrow \nu t \bar{b} X$. However, the production cross-section for this process at HERA energy is very small ($\sigma(ep \rightarrow tX) < 1$ fb [12]) and far beyond an experimental observation. On the other hand the anomalous FCNC process could contribute to the single top production at HERA energies. Therefore, a study of such reaction at HERA provide a good possibility to put a constraint on FCNC interactions.

The both H1 and ZEUS collaborations searched for the top quark production in reaction (1). No evidence of the top production was found. The resulted constraint on the single top production cross-section is given below [8, 9]:

$$\sigma(ep \rightarrow etX) < 0.225 \text{ pb} \quad (\text{at } 95\% \text{ CL}) \quad (2)$$

The conventional interpretation of this data assumes that the top quark anomalous FCNC interactions could occur with a photon or Z -boson only (see Fig. 1). As a result from

evaluated [8, 9]. In terms of BR this constraint is as follows:

$$\text{BR}(t \rightarrow q\gamma) < 0.7\% \quad (3)$$

In the present article we consider an additional FCNC process that contribute to the single top-quark production at HERA energies. Indeed, in ep -collisions the single top could be produced also due to anomalous interaction with a gluon, i.e. tgq vertex, as it is shown on fig. 2. Thereby using the upper limit on the production cross-section (2) we evaluate a constraint on anomalous FCNC coupling of the top quark with a gluon. Therein lies the basic idea of our analysis.

1 Phenomenological Lagrangian for FCNC

We do not know which type of new physics will be responsible for a future deviation from the SM predictions. However, the top quark couplings can be parameterized in a model independent way by an effective Lagrangian [1] (see also [13] for the first discussion of the $t\bar{c}g$ -operator).

In our article we consider the top quark anomalous interactions with gluon and photon. However for the completeness we also present here the expression for the t -quark FCNC interaction with Z -boson. Actually the width of $t \rightarrow Zq$ decay is used in the Section 4. Thus the effective phenomenological Lagrangian have a form as follows:

$$\begin{aligned} \mathcal{L} = & - 2g_s \sum \frac{\kappa_g}{\Lambda} \bar{t} \sigma_{\mu\nu} T^a (g_L P_L + g_R P_R) u q^\nu G_a^\mu \\ & - 2e \sum \frac{\kappa_\gamma}{\Lambda} \bar{t} \sigma_{\mu\nu} (\gamma_L P_L + \gamma_R P_R) u q^\nu A^\mu \\ & - \frac{e}{\sin 2\vartheta_W} \sum \kappa_Z \bar{t} \gamma^\mu (z_L P_L + z_R P_R) u Z_\mu + h.c., \end{aligned} \quad (4)$$

where t^a are the Gell-Mann matrices satisfying $\text{Tr}(t^a t^b) = \delta^{ab}/2$; κ_g and κ_γ define the strength of the FCNC with a gluon and a photon respectively (real and positive values); $\Lambda = 1$ TeV is a new physics cutoff; e is the electric charge; ϑ_W is the Weinberg angle; $\sigma^{\mu\nu} = \frac{1}{2}(\gamma^\mu \gamma^\nu - \gamma^\nu \gamma^\mu)$; $P_{R,L} = \frac{1}{2}(1 \pm \gamma^5)$; $g_L^2 + g_R^2 = \gamma_L^2 + \gamma_R^2 = 1$; q^ν is the momentum of corresponding gauge boson.

The top-quark decay widths into one of the light quarks (u or c) with a gluon, photon and Z -boson have the form:

$$\Gamma(t \rightarrow gq) = \frac{4}{3} \alpha_s m_t^3 \left(\frac{\kappa_g}{\Lambda} \right)^2 \quad (5)$$

$$\Gamma(t \rightarrow \gamma q) = \alpha m_t^3 \left(\frac{\kappa_\gamma}{\Lambda} \right)^2 \quad (6)$$

$$\Gamma(t \rightarrow Zq) = \frac{\alpha \kappa_Z^2 m_t^3}{8 \sin^2 2\vartheta_W M_Z^2} \left(1 - \frac{M_Z^2}{m_t^2} \right)^2 \left(1 + 2 \frac{M_Z^2}{m_t^2} \right) \quad (7)$$

Firstly, we re-calculate the matrix element squared for $2 \rightarrow 2$ process: $eu \rightarrow et$ (see the diagram on Fig. 1). This expression is used in our analysis. The explicit form [15] is given below:

$$|M|^2 = 4e^4 \left(\frac{1}{Q^2} \right)^2 \left(\frac{\kappa_\gamma}{\Lambda} \right)^2 \times \{ 2m_e^2 \hat{t}(2\hat{s} + \hat{t}) - 2m_e^2 m_t^4 - \hat{t}[2\hat{s}(\hat{s} - m_t^2) + m_t^4] - \hat{t}^2(2\hat{s} - m_t^2) \} \quad (8)$$

where $\hat{s}$ and $\hat{t}$ are usual Mandelstam variables, $Q^2 = |\hat{t}|$ is a photon transferred momentum squared, m_e and m_t are the masses of the electron and t -quark. Note, that we keep non-zero electron mass, because this is essential for the further analysis.

One can obtain an expression for the production cross-section of the hard $2 \rightarrow 2$ process (corresponding to matrix element squared (8)) by the integration on Q^2 from Q_{min}^2 to Q_{max}^2 :

$$\begin{aligned} \hat{\sigma}(Q_{min}^2) &= C \left[B \ln \left(\frac{Q_{max}^2}{Q_{min}^2} \right) - A(Q_{max}^2 - Q_{min}^2) \right] \\ &- 2Cm_e^2 \left[2\hat{s} \ln \left(\frac{Q_{max}^2}{Q_{min}^2} \right) - (Q_{max}^2 - Q_{min}^2) \right] \\ &+ 2Cm_e^2 m_t^4 \left(\frac{1}{Q_{min}^2} - \frac{1}{Q_{max}^2} \right), \end{aligned} \quad (9)$$

where

$$C = \frac{\pi e^4}{2\hat{s}(\hat{s} - m_e^2)} \left(\frac{\kappa_\gamma}{\Lambda} \right)^2, \quad A = 2\hat{s} - m_t^2, \quad B = 2\hat{s}(\hat{s} - m_t^2) + m_t^4$$

The single top production in ep -collisions due to anomalous FCNC interaction with a gluon is described by two subprocesses with a gluon or $u(c)$ -quark from the initial proton (see the diagrams on Fig. 2):

$$eg \rightarrow etq, \quad q = u \text{ or } c \quad (10)$$

$$eq \rightarrow etg, \quad q = u \text{ or } c \quad (11)$$

Note, that the matrix elements squared for these two processes are related to each other by crossing. Therefore, we present an exact expression for $|M|^2$ for the first subprocess (10):

$$|T|^2 = |T_1 + T_2|^2 \quad (12)$$

The exact expressions are given below:

$$\begin{aligned} |T_1|^2 &= A_0 A_1^2(kp_2) \left\{ (f_1 l_1)(p_1 l_2) + (f_1 l_2)(p_1 l_1) - (p_1 f_1) \left[(l_1 l_2) + \frac{q^2}{2} \right] \right\} \\ &- 2A_0 A_1^2 m_t^2(kp_2) [(l_1 l_2) + q^2]; \\ |T_2|^2 &= A_0 A_2^2(kp_1) \left\{ (f_2 l_1)(p_2 l_2) + (f_2 l_2)(p_2 l_1) - (p_2 f_2) \left[(l_1 l_2) + \frac{q^2}{2} \right] \right\}; \end{aligned}$$

$$\begin{aligned}
\frac{T_1^+ T_2 + T_2^+ T_1}{A_0 A_1 A_2} &= 4(p_1 l_1)(p_1 l_2)(k p_2)^2 + 4(p_2 l_1)(p_2 l_2)(k p_1)^2 + 4(k l_1)(k l_2)(p_1 p_2)^2 \\
&- 2(p_1 p_2) \{2(f_3 l_1)(k l_2) + 2(f_3 l_2)(k l_1) - (k f_3)(l_1 l_2)\} \\
&+ 2m_t^2(k p_2) \{(p_2 l_1)(k l_2) + (p_2 l_2)(k l_1) - (k p_2)(l_1 l_2)\} \\
&- m_t^2 \{2(f_3 l_1)(k l_2) + 2(f_3 l_2)(k l_1) - (k f_3)q^2\} \\
&+ 2m_t^2(k l_1)(k l_2)(p_1 p_2);
\end{aligned}$$

where

$$A_0 = 64 \left[\frac{4}{3} \frac{\kappa_g}{\Lambda} \alpha_s e^2 \right]^2, \quad A_1 = -\frac{1}{2(k p_2) + m_t^2}, \quad A_2 = -\frac{1}{2(k p_1) - m_t^2}$$

$$f_1 = 2(k p_2)p_2 + m_t^2 k, \quad f_2 = 2(k p_1)p_1 - m_t^2 k, \quad f_3 = (k p_1)p_2 + (k p_2)p_1.$$

here $l_1(l_2)$ are 4-momentum of the initial (final) electron, p_1 and p_2 are 4-momenta of the top- and $\bar{u}$ -quarks, and k is the initial gluon momentum. The calculation of the color coefficients is trivial and they are included to A_0 .

For numerical calculations the evaluated matrix elements squared are incorporated in the event generator TopReX [14].

3 Kinematics of the considered processes

The cross-section of the considered processes depends essentially on the kinematical limit of the gauge boson transferred momentum Q_{min} . This limit is determined by the additional requirements on kinematics of the reaction and the detector parameters. For some reason the authors of the experimental paper [9] did not mention the limit on Q_{min} in an explicit form.

In order to calculate the theoretical value of the cross-section, σ^{th} , we must imply a correct limit on Q_{min} . This value can be estimated from the following considerations. The experimental analysis for $2 \rightarrow 2$ process was made under the assumption that single top is produced due to anomalous coupling with a photon. It is reasonable that the theoretical cross-section also depends on the Q_{min} kinematical limit and anomalous coupling with a photon. Therefore by using the value of the anomalous coupling one can compare theoretical cross-section with experimental one and can determine the unknown value of Q_{min}^2 .

Thus we get the value of Q_{min}^2 from the following condition:

$$\sigma_{2 \rightarrow 2}^{\text{th}}(ep \rightarrow etX) = \left(\frac{\kappa_\gamma}{\Lambda} \right)^2 \times f_{2 \rightarrow 2}^{ep \rightarrow etX}(Q_{min}^2) \leq \sigma^{exp} = 0.225 \text{ pb} \quad (13)$$

Here $\sigma_{2 \rightarrow 2}^{\text{th}}$ is the total cross-section for $ep \rightarrow etX$ process (with subprocess $eu \rightarrow et$) calculated using the expression (9), σ^{exp} is the experimental limit on the single top quark production from the HERA data [9]. The upper limit on anomalous coupling of the top quark with a photon was taken from the same article [9]: $\kappa_{tu\gamma}^H < 0.174$.

phenomenological Lagrangian:

$$\mathcal{L}^{\mathcal{H}} = ee_t \bar{t} \frac{\sigma_{\mu\nu} q^\nu}{\Lambda^H} u \kappa_{tu\gamma}^H,$$

where e is the electric charge, e_t is the charge of up quarks and $\Lambda^H = m_t = 175$ GeV.

The relation between $\kappa_{tu\gamma}^H$ and κ_γ used in our Lagrangian (4) looks as follows:

$$\kappa_\gamma = \frac{e_t}{\sqrt{2}} \frac{\Lambda}{\Lambda^H} \frac{1}{\sqrt{\gamma_L^2 + \gamma_R^2}} \kappa_{tu\gamma}^H \quad (14)$$

Note that we take for the new physics cutoff $\Lambda = 1$ TeV and by our convention $\gamma_L^2 + \gamma_R^2 = 1$ (see Eq. (4)). As a result the upper limit on the anomalous coupling in terms of the Lagrangian (4) equals:

$$\kappa_\gamma = 0.468. \quad (15)$$

The single top production cross-section in ep -collisions was evaluated within the parton model:

$$\sigma^{\text{th}}(e^\pm p \rightarrow e^\pm t X) = \int_{m_t^2/s}^1 dx f(x, \mu^2) \hat{\sigma}(x, \mu^2), \quad (16)$$

where x is the part of the proton momentum carried away by a parton; $\hat{\sigma}(x, \mu^2)$ is the cross-section for the hard subprocess; $f(x, \mu^2)$ is the parton distribution function, and $\mu = m_t$ is the factorization scale.

Using (13) we obtain

$$Q_{min}^2 \geq 0.08 \text{ GeV}^2 \quad (17)$$

Further in our analysis of the single top production due to anomalous coupling with a **gluon** we implied the very same limit on Q_{min}^2 (17).

4 The constraint on FCNC coupling of the top-quark with a gluon

In the present study we compare the experimental constraint (2) with the total top-quark production cross-section in ep -collisions (evaluated with the matrix elements squared (12)).

$$ep \rightarrow et f X, \quad (18)$$

Thus the single top quark production in ep -collisions is possible not only via FCNC with a photon, but also due to anomalous interactions with a gluon. This idea gives us an opportunity to obtain the direct constraint on FCNC coupling of the top quark with a gluon from HERA experimental data (2).

In order to find the constraint on gluon coupling κ_g we use the following relation:

$$\sigma^{th}(ep \rightarrow etfX) \times BR(t \rightarrow bW^+) \leq \sigma^{exp} = 0.225 \text{ pb}, \quad (19)$$

where

$$\sigma_{2 \rightarrow 3}^{th}(ep \rightarrow etfX) = \left(\frac{\kappa_g}{\Lambda}\right)^2 \times F_{2 \rightarrow 3}^{ep \rightarrow etfX}(Q_{min}^2).$$

Here we take into account the possible modification of the branching ratio of the top decay in the SM mode $t \rightarrow bW^+$. As it was mentioned above the top quark anomalous interactions could lead to the appearance of additional top decay channels. So the probability of the SM top decay would change accordingly:

$$BR(t \rightarrow bW^+) = \frac{\Gamma(t \rightarrow bW^+)}{\Gamma(t \rightarrow bW^+) + \Gamma(t \rightarrow Zq) + \Gamma(t \rightarrow gq)}, \quad (20)$$

To obtain an upper limit on the anomalous coupling with a gluon we assume the absence of the top-quark FCNC interaction with a photon, i.e. $\kappa_{t\gamma u,c} = 0$. In addition the different cases of the top interactions with a gluon and Z -boson are explored:

1. $\kappa_{tZu,c} = 0, \kappa_{tgc} = 0, \kappa_{tgu} \neq 0$,
the simplest case, i.e. gluon - top - u -quark interaction;
2. $\kappa_{tZu,c} = 0, \kappa_{tgc} = \kappa_{tgu} \neq 0$.
here the FCNC transitions with a gluon to u - or c -quark are considered;

Two additional cases include the anomalous interaction with Z -boson:

3. $\kappa_{tZu,c} \neq 0, \kappa_{tgc} = 0, \kappa_{tgu} \neq 0$.
4. $\kappa_{tZu,c} \neq 0, \kappa_{tgc} = \kappa_{tgu} \neq 0$.

According to all assumptions (1.-4.) the constraints on the top quark FCNC couplings with a gluon, u -, c -quarks and the corresponding BR of the anomalous top decays are calculated. The results of the analysis are presented in Table 2.

Table 2: *The constraints on the top quark FCNC coupling with a gluon from the HERA data ($\kappa_{t\gamma q} = 0$)*

		$\kappa_{tgu}/\Lambda, \text{ TeV}^{-1}$	$\kappa_{tgc}/\Lambda, \text{ TeV}^{-1}$	$BR(t \rightarrow gq)$
1	$\kappa_{tZq} = 0, \kappa_{tgc} = 0$	0.393	—	6.6%
2	$\kappa_{tZq} = 0, \kappa_{tgu} = \kappa_{tgc}$	0.396	0.396	12.6%
3	$\kappa_{tZq} \neq 0, \kappa_{tgc} = 0$	0.408	—	6.2%
4	$\kappa_{tZq} \neq 0, \kappa_{tgu} = \kappa_{tgc}$	0.410	0.410	11.8%

The best constraint on FCNC coupling with a gluon from ep -collision is obtained supposing the interaction with an u -quark only. When the interaction with a charm quark is also

almost twice. It is explained by the small contribution of the c -quarks to proton's PDF and appearance of two decay modes of the top-quark. One also can see that the contribution of Z -boson modifies the results slightly and can be neglected.

Finally the derived constraint on the anomalous coupling of the top FCNC interaction with a gluon and the corresponding BR are given below:

$$|\kappa_{tgq}/\Lambda| \lesssim 0.4 \text{ TeV}^{-1}, \quad \text{BR}(t \rightarrow gq) < 13\% \quad (21)$$

5 Conclusions

Very recently at the Tevatron experiment [11] better constraint ($3 \div 10$ times less) on the κ_g , the upper limit on the anomalous coupling of the top quark with a gluon, was obtained.

Nevertheless we wish to indicate that the HERA results for the single top production provides the way to put the constraints on the anomalous top-quark interactions not only with a photon and Z -boson, but also with a gluon!

Moreover, such analysis of ep -collision (e.g. HERA data) allows getting the limits on anomalous coupling of the top quark with a gluon and can be considered as an independent estimation of κ_g .

Acknowledgments

We are grateful to M. Mohammadi Najafabadi and O.V. Zenin for useful discussions. This work was supported in part by the Dynasty Foundation and by the Russian Foundation for Basic Research under Grant # 08-02-91002-C.

References

- [1] Beneke M. *et al.*, “*Top quark physics*”, in “*Standard model physics (and more) at the LHC*”, G. Altarelli and M. L. Mangano eds., *Geneva, Switzerland: CERN (2000)* [arXiv:hep-ph/0003033].
- [2] Grzadkowski B., Gunion J.F., and Krawczyk P., *Phys. Lett.*, **B268**, 106 (1991);
Eilam G., Hewett J.L, and Soni A., *Phys. Rev.* **D44**, 1473 (1991);
Luke M. and Savage M.J., *Phys. Lett.* **B307**, 387 (1993);
Couture G., Hamzaoui C., and K  nig H., *Phys. Rev.* **D52**, 1713 (1995).
- [3] Atwood D., Reina L. and Soni A., *Phys. Rev.*, **D55**, 3156 (1997) [arXiv:hep-ph/9609279].
- [4] Yang J.M., Young B.-L. and Zhang X., *Phys. Rev.*, **D58**, 055001 (1998) [arXiv:hep-ph/9705341].

- [5] [arXiv:hep-ph/9808400].
- [6] Abe F. *et al.* [CDF Collaboration], *Phys. Rev. Lett.* **80**, 2525 (1998).
- [7] Wolf G., [arXiv:hep-ex/0105055];
 Alan A.T. and Senol A., *Europhys. Lett.* **59**, 669 (2002) [arXiv:hep-ph/0202119];
 Dannheim H. [H1 Collaboration], [arXiv:hep-ex/0212004].
- [8] Aktas A. *et al.* [H1 Collaboration], *Eur. Phys. J.* **C33**, 9 (2004) [arXiv:hep-ex/0310032].
- [9] Chekanov S. *et al.* [ZEUS Collaboration], *Phys. Lett.* **B559**, 153 (2003) [arXiv:hep-ex/0302010].
- [10] Obraztsov V.F., Slabospitsky S.R., and Yushchenko O.P., *Phys. Lett.* **B426**, 393 (1998) [arXiv:hep-ph/9712394];
 Heister A. *et al.* [ALEPH Collaboration], *Phys. Lett.* **B543**, 173 (2002) [arXiv:hep-ex/0206070];
 Abreu P. *et al.* [DELPHI Collaboration], *Phys. Lett.* **B446**, 62 (1999) [arXiv:hep-ex/9903072];
 Abdallah J. *et al.* [DELPHI Collaboration], *Phys. Lett.* **B590**, 21 (2004) [arXiv:hep-ex/0404014];
 Achard P. *et al.* [L3 Collaboration], *Phys. Lett.* **B549**, 290 (2002) [arXiv:hep-ex/0210041];
 Abbiendi G. *et al.* [OPAL Collaboration], *Phys. Lett.* **B521**, 181 (2001) [arXiv:hep-ex/0110009].
- [11] Abazov V.M. *et al.* [D0 Collaboration] *Phys. Lett.* **99**, 191802 (2007) [arXiv:hep-ex/0702005].
- [12] Stelzer T., Sullivan Z. and Willenbrock S., *Phys. Rev.* **D 56**, 5919 (1997) [arXiv:hep-ph/9705398];
 Moretti S. and Odagiri K., *Phys. Rev.* **D 57**, 3040 (1998) [arXiv:hep-ph/9709435].
- [13] Malkawi E. and Tait T., *Phys. Rev.* **D54** (1996) 5758 [arXiv:hep-ph/9511337]
 Tait T. and Yuan C. P., *Phys. Rev.* **D55** (1997) 7300 [arXiv:hep-ph/9611244].
- [14] Slabospitsky S.R. and Sonnenschein L., "TopReX generator (version 3.25): Short manual", *Comput. Phys. Commun.* **148**, 87 (2002) [arXiv:hep-ph/0201292].
- [15] Belyaev A. and Kidonakis N., *Phys. Rev.* **D65**, 037501 (2002).




Figure 1: *Single top production via FCNC coupling with photon (or Z-boson)*




Figure 2: *The Feynman diagrams of the single top production in ep-collisions due to FCNC with a gluon*